

Canadian Translation of Fisheries and Aquatic Sciences

No. 5188

The relation between sediment flux and fish feeding
in coastal culture farms

S. Kadowaki, T. Kasedo, T. Nakazono,
Y. Yamashita, and H. Hirata

Original title: Senkai Yoshoku Cyozo ni okeru Chinkosei Busshitsu to
Kyujiryo narabi Busshitsu to no Kankei

In: Mem. Fac. Fish. Kagoshima Univ. 29: 217-224, 1980

Original language: Japanese

Available from:

Canada Institute for Scientific and Technical Information
National Research Council
Ottawa, Ontario, Canada K1A 0S2

1985

13 typescript pages

Fisheries & Oceans
LIBRARY

JAN 15 1987

BIBLIOTHÈQUE
Pêches & Océans



Secretary
of State

Secrétariat
d'État

90-00071/

MULTILINGUAL SERVICES DIVISION - DIVISION DES SERVICES MULTILINGUES

TRANSLATION BUREAU

BUREAU DES TRADUCTIONS

LIBRARY IDENTIFICATION - FICHE SIGNALÉTIQUE

CTFAS 5188

Translated from - Traduction de Japanese	Into - En English
Author - Auteur	

Shusaku KADOWAKI, Teruo KASEDO, Tsurayuki NAKAZONO, Yaoki YAMASHITA and Hachiro HIRATA

Title in English or French - Titre anglais ou français

The Relation between Sediment Flux and Fish Feeding in Coastal Culture Farms

Title in foreign language (Transliterate foreign characters)
Titre en langue étrangère (Transcrire en caractères romains)

Senkai Yoshoku Cyoyo ni okeru Chinkosei Busshitsu to Kyujiryo narabini Jishitsu
to no Kankei

Reference in foreign language (Name of book or publication) in full, transliterate foreign characters.
Référence en langue étrangère (Nom du livre ou publication), au complet, transcrire en caractères romains.

Kagoshima Daigaku Suisanbu Kiyo

Reference in English or French - Référence en anglais ou français

Mem. Fac. Fish. Kagoshima Univ.

Publisher - Editeur	DATE OF PUBLICATION DATE DE PUBLICATION			Page Numbers in original Numéros des pages dans l'original
	Year Année	Volume	Issue No. Numéro	217-224
Piece of Publication Lieu de publication				Number of typed pages Nombre de pages dactylographiées
Japan	1980	29	-	13

Requesting Department
Ministère-Client **DFO**

Translation Bureau No.
Notre dossier n° **2126513**

Branch or Division
Direction ou Division **SIPB**

Translation (Initials)
Traducteur (Initiales) **P.S.**

Person requesting
Demandé par **A. T. Reid**

Your Number
Votre dossier n° **-**

Date of Request
Date de la demande **September 10, 1985**

OCT 30 1985



MULTILINGUAL SERVICES DIVISION — DIVISION DES SERVICES MULTILINGUES

TRANSLATION BUREAU

BUREAU DES TRADUCTIONS

Client's No.—N ^o du client	Department — Ministère	Division/Branch — Division/Direction	City — Ville
	DFO	SIPB	Ottawa
Bureau No.—N ^o du bureau	Language — Langue	Translator (Initials) — Traducteur (Initiales)	
2126513	Japanese	P.S.	OCT 30 1985

The Relation between Sediment Flux and Fish Feeding
in Coastal Culture Farms*¹Shusaku KADOWAKI*², Teruo KASEDO*², Tsurayuki NAKAZONO*²,
Yaoki YAMASHITA*³ and Hachiro HIRATA*⁴

Abstract

Sediment flux in coastal culture farms was observed in order to estimate an environmental load resulting from fish feeding. Observations were made 63 times, between October 1977 and February 1980, in the farms located around the Fisheries Research Laboratory of Kagoshima University.

The sediments were collected from 3 stations: St. 1 in a sea bream farm, St. 2 in a yellow-tail farm, and St. 3 in a non-culture area. Water depth in the St. 1, 2, and 3 were: 8 to 12 m, 14 to 16 m, and 12 to 13 m, respectively. Polyethylene columns, 5 liters capacity with 17 cm opening, were set on the sea bottom at each station, and renewed every 4-days during the culture seasons. The samples were filtered after collection, and dried at 60°C for 5 days. The food quantity and quality were noted every day in each farm.

Average daily amounts of sediments in St. 1, 2, and 3 were calculated to be: 33.8, 23.0, and 14.6 g/m², respectively. The sedimentations were related to the amount of foods supplied, and to water depth. Some diffusion of biodeposits was observed in a deeper farm. The type of food supplied also affected the amount of sediment flux. The amounts of sediments ranged from 4.1 to 5.9 g/m² when the fish were fed daily on 1202 kg of fresh anchovies, but such amounts increased about 4 times, 17.0 to 21.6 g/m², when 1247 kg of frozen mackerel were supplied.

*¹ (Contribution No. 14 from Fish. Res. Lab., Fac. Fish., Kagoshima Univ.).

*² (Fish. Res. Lab., Fac. Fish., Kagoshima Univ., Azuma-cho, Izumi-gun, Kagoshima, 899-14 Japan).

*³ (Yamashita Fish. Co. Ltd., Azuma-cho, Izumi-gun, Kagoshima, 899-14 Japan).

*⁴ (Lab. Fish Cultivation Physiol., Fac. Fish., Kagoshima Univ., Kagoshima, 890 Japan).

UNEDITED TRANSLATION
For information only
TRADUCTION NON REVISEE
Information seulement

UNEDITED TRANSLATION
For information only
TRANSDUCTION NON REVISEE
Information seulement

2

Foreword

p. 218

Research about fish farming in shallow waters and auto-pollution was taken as a topic at the 1977 spring convention of the Japanese Society for Scientific Fisheries. We see from the collected lectures that many reports have already been produced. However, about the main topic which is the relationship between the volume of sediment and the volume of feed as well as the feed quality, the analyses are still insufficient. Regarding the relationship between the amounts of feed and volumes of sediment, Hagino (1977) compiled the results of tests conducted in tanks at the Wakayama and at the Kochi Fisheries research centers. The data from each center are, however, still few.

The main reason for the few data about the volumes of feed and of sediment is that it is extremely difficult to gather data at the farming grounds. More particularly, the difficulty of knowing the precise amounts of feeding at the site is a well known fact. Fortunately a large-scale shallow farming ground is extending in front of this fisheries research center and since 1977 it has been possible to the farm operators and to this research center staff to carry out research in common. We were therefore able to establish the correlation between the volumes of sediment, of feeding and the feed quality.

As research on sediments accompanying fish farming is a relatively recent field, the terminology is still unsettled (Hagino 1977; Hata 1979, Koyama 1975, Tanaka 1977). So we decided to call "sediment flux" the volume of sediment precipitation for one day per unit area, and expressed its unit as $\text{g}/\text{m}^2 \cdot \text{day}$.

The present study was commissioned by the Azuma town's council, the Kagoshima Prefecture, by the Azuma Fisheries Cooperative and by

the National Seawater Fish Farming Association. We thank all those involved for their support. We also thank Professor Akira FUJI of the Hokkaido University and Dr. Keiyo TANAKA of the Kagawa University for their kind advice. We express also our thanks to Akihiko SHINOMIYA, instructor at this department, and to Mr. Ko YAMASHITA of the Yamashita Fisheries Co. for their kind cooperation.

Survey method

According to the classification made by INOUE et al (1970), the presently surveyed fishing ground is a typical fish farm using a small channel. The variations of the tide moving back and forth between the East China Sea and Yatsushiro Sea are strong and the current in the vicinity of the fish cages reaches often 1-2 kt. As shown in Fig. 1, the area of the present fish farm is roughly 10 ha, and about 300 cages (mainly $7 \times 7 \times 5 \text{ m}^3$) are floating at this site. The annual production of this fish farm is estimated at more or less 400 t.

We surveyed three sites viz. a cluster of cages for the rearing of red seabream (St. 1; a-c), a cluster of cages for the rearing of yellowtail (St. 2; a-c) as well as a non-farming fishing ground located about 500 m away from the above cage clusters (St. 3; a or b). The depth at St. 1 is 8-12 m and thus relatively small. At St. 2 and St. 3, the depths are 14-16 m and 12-13 m respectively. As for the survey of St. 3, we operated at the point a until December 1977, and thereafter at the point b (Fig. 1).

To gather the sediments, we followed the method of Kadowaki et al (1978). That is, we suspended 25 cm long cylindrical polyethylene containers with a 17-cm opening by means of floaters and weights 60-80 cm from the sea floor, collected almost every 4 days the sediment gathered

inside the container, filtered the sediment with No. 2 filter paper, and weighed carefully after heating for 2 days at 60°C. The feeding and the quality of the feed were recorded daily for each cage group during the survey period.

From October 1977 to February 1980, the survey was conducted intermittently. In all, we operated 63 times on 285 days. From March to May in the spring, the fish cages are replaced and moved so that no survey was possible during that time. For 3 years we performed regular surveys every four days during the maximum feeding period of October-December.

p. 219

UNEDITED TRANSLATION

For information only

TRADUCTION NON REVISEE

Information seulement

Results and discussion

Table 1 shows all the data of the surveys. On the basis of Table 1, we found the respective 3-point displacement average values of the feeding volume and the sediment flux. In Fig. 2 we showed these correlations by St. (by water depth). Although the relationship between the volumes of feed and the sediment flux shows scattering for all stations, we saw a rough positive correlation in all cases (at St. 1, $r = +0.49$ and at St. 2, $r = +0.58$).

As the Fig. shows, the slope at St. 1 (red seabream cage cluster) is relatively steep while at St. 2 (yellowtail cages), the slope tends to be gentle. Rather than being due to the species of fish, the difference is due to the fact that while the water depth at St. 1 is rather small with 8-12 m, it is 12-16 m at St. 2 and there seems to be a wider dispersion of suspended matters where the depth under the cage cluster is greater.

p. 223

As indicated in the bottom column of Table 2 which gives the sediment flux per 1 t of feeding, a-c of St. 1 shows always an average

value of 28.9 dry $\text{g}/\text{m}^2 \cdot \text{day}$. In a-c of St. 2, the average was 13.6 dry $\text{g}/\text{m}^2 \cdot \text{day}$. If basing ourselves on the instance of rotifer rearing by Yamauchi (1979), we make the dry-wet conversion, we obtain 444.5 wet $\text{g}/\text{m}^2 \cdot \text{day}$ for St. 1 and 209.2 wet $\text{g}/\text{m}^2 \cdot \text{day}$ for St. 2. So, the rates of sediment flux (wet weight) per 1 t of feeding in St. 1 and in St. 2 are about 30 and 15% respectively.

Hagino (1977) states that the feeding volume-sediment flux ratio in mid-bay fish farms is about 10%. However, in the channel-type fish farm such as the ones studied in this survey, relatively higher values were obtained. These differences seem to be due to differences in dispersion concomitant to such conditions as depth.

Meanwhile the sediment flux at the non-rearing fishing ground St. 3 removed about 500 m from the fish cage clusters showed an average of 14.6 dry $\text{g}/\text{m}^2 \cdot \text{day}$ throughout the entire period. This value is about half of the values obtained at the fish farms of St. 1 and St. 2 (Table 1). These facts mean a dispersion of floating and suspended matters from the fish farms to the non-farming fishing grounds. One should make additional surveys taking into account factors such as depth and currents.

Table 2 shows the seasonal variations of the sediment flux at the fish farms. The average is highest in the fall from September to November, with 36.7 dry $\text{g}/\text{m}^2 \cdot \text{day}$. For the winter (Dec.-Jan.) and the summer (June-July), the average is 45% lower than in the fall. The average was 16.4 dry $\text{g}/\text{m}^2 \cdot \text{day}$. The observed high sediment flux in the fall seems to correspond to the maximum feeding taking place at that time.

One noteworthy point is that the sediment flux depends on the type of feed. As Table 1 shows, frozen mackerel was given from June 6

UNEDITED TRANSLATION
For information only
TRADUCTION NON REVISEE
Information seulement

6

p. 224

to June 26, 1978. From the following 27th to July 13, the feed consisted in fresh anchovies which had just been caught. The difference in sediment fluxes between these 2 types of food is obvious and we compiled the average values in Table 3. The amounts of both feeds per day were about equal at 1,200-1,300 kg/day. While the average sediment flux was 17.0-21.6 dry $\text{g/m}^2 \cdot \text{day}$ during the frozen mackerel feeding period, it was only 4.1-5.9 dry $\text{g/m}^2 \cdot \text{day}$ viz. $\frac{1}{4}$ this volume when feeding the fish fresh anchovies. The volume was thus extremely small. These fresh anchovies had been caught by the operators and one may say that the difference in feed quality and excellent freshness contributed to bring down the sediment flux. One might consequently appreciate the fact that if fish farmers can secure feed by themselves, ideal conditions for fish farming in shallow waters will be obtained.

Recent studies about sediments in the sea are not merely related to fish farming (Nitta 1980, Tanaka 1977). We notice new topics in the fields of marine physics and chemistry (Handa 1979; Hata 1979; Hirata et al 1979; Iseki et al 1980; Nishizawa 1966).

Handa wrote comprehensively about sediment traps (1979) and clarified various phenomena regarding sediments in the sea. Meanwhile, auto-pollution accompanying fish farming is studied carefully in connection with the red tide problem (Hanaoka, 1980; Nitta, 1980; Tahira, 1978). Thus the importance of research about bottom sediments in the sea increases from both the viewpoints of natural and of social sciences. In the future we wish to pursue our surveys of the bottom sedimentation mechanism and of the state of dispersion at the fishing grounds while referring ourselves to the present results and to the continuous DO recording data by cruising in fish farming grounds.




Fig. 1. Map showing the experimental stations in a coastal fish farm with isobath.

Table 1. Daily amount of sediment flux and food supplied.

Item	SEDIMENT (dry g/m ² ·day)							FOOD (kg dry)	
Farm	Sea bream (St. 1)			Yellowtail (St. 2)			Out of farm	Sea bream	Yellow-tail
Substation Depth	a 8 m	b 10 m	c 12 m	a 14 m	b 15 m	c 16 m	(St. 3 a or b) 12~13 m		
1977									
Oct. 13-17		47.5		23.2	61.9	67.2	36.5	1770	1120
17-21				32.1	59.7	39.8	21.0	1010	2240
21-25				44.2	58.6	49.7	24.3	760	1370
25-29		51.9		45.3	84.0	32.1	18.8	760	1370
29-2		65.2		33.2	22.1	32.1	17.7	1590	1360
Nov. 2-7		59.7		44.2	19.9	24.3	24.3	3850	2430
7-11		47.5		51.9	32.1	26.5	30.9	1010	1530
11-15		40.9		47.5	24.3	28.7	37.1	840	2970
15-19		36.5		34.3	26.5	29.8	21.0	1460	1440
19-24		22.1		45.3	18.8	19.9	14.4	2080	1150
24-28		16.6		51.9	25.4	28.7	16.6	1390	2870
28-2		18.8		37.6	18.8	11.1	15.5	1130	1260
Dec. 2-7		19.5		32.1	13.3	12.2	11.7	540	920
7-12		23.2		19.9	14.4	12.2	13.9	1020	430
12-16		21.0		18.8	17.7	14.4	19.4	530	1670
1978									
June 6-10	11.4	19.5	9.5	18.9	15.5	14.3	16.5	980	850
10-14	20.8	21.3	15.3	10.6	9.2	6.0	5.8	1950	1030
14-18	22.7	48.0	45.3	19.6	23.2	27.9	9.5	750	1090
18-22	20.8	32.4	21.0	21.4	20.0	19.8	15.8	1240	2360
22-26	13.7	12.0	10.7	14.7	17.1	17.2	20.1	980	1240
26-1	3.1	6.8	2.2	4.9	5.3	4.6	4.1	1510	1190
July 1-6	8.4	9.1	8.3	3.9	2.6		6.0	1560	720
9-13	3.5	4.3	7.7	3.3				990	1240
Sep. 5-9	55.1	53.0	25.4				13.1	1050	
9-13	68.4	32.6	35.3				11.8	1660	
13-17	50.2	58.5	49.9				28.4	710	
17-21	72.3	30.3	17.8				28.8	1500	
21-25	42.5	52.6	12.2				23.0	1690	
25-29	21.5	49.8	38.9				6.4	1800	
Oct. 3-7	57.6	93.3	53.1				28.1	1910	
7-11	63.9	60.5	44.5				24.4	940	
11-15	60.7	59.8	50.5				16.8	940	
15-19	44.1	37.6	41.7					1500	

Table 1. (Continued)

Item		SEDIMENT (<i>dry-g/m²-day</i>)						FOOD (<i>kg/day</i>)	
Farm		Sea bream (St. 1)			Yellowtail (St. 2)			Out of farm	
Substation	Depth	a	b	c	a	b	c	(St. 3 a or b) 12~13 m	Sea bream Yellowtail
		8 m	10 m	12 m	14 m	15 m	16 m		
	19-23	38.0	54.0	31.2					1760
	23-27	42.8	59.1	23.4				3.4	1860
	27-31	25.4	70.7	42.7				19.6	1900
	31-4	36.8	46.3	43.8				15.7	1800
Nov.	4-8	33.4	57.8					6.7	870
	12-16	22.2	42.1	64.3				5.5	1880
	16-20	28.0	20.4	16.8				4.3	900
	20-24	44.8						10.3	940
	24-28	50.4	23.9	27.9				5.6	640
	28-2	71.0	27.1	23.2				12.9	1120
Dec.	2-7	70.9	21.8	11.6				10.8	730
	7-11	30.5	30.0	14.8				9.3	480
	14-18	11.8	17.3	7.0				6.9	560
1979									
Oct.	8-12	31.5	49.1	46.8	13.1	9.6	4.4	5.1	580
	12-16	31.5	31.0	22.2	41.4	39.8	22.6	6.3	1000
	16-20	55.1	67.7	41.7	66.2	8.5	11.4	9.9	540
	20-24	21.6	60.9	54.7	21.5	5.1	6.5	12.1	1200
	28-1	19.3	101.7	17.6	3.6	12.8	11.2	9.8	990
Nov.	1-5	12.1	51.9	52.5	19.5	11.7	11.4	11.4	480
	5-9	19.9	67.4	43.5	22.9	9.4	18.8	17.8	1020
	9-13	8.5	135.9	29.5	22.8	29.2	12.3	11.0	1110
	13-17	21.8	53.6	43.7	9.8	41.3	15.5	16.8	1050
	17-22	18.6	150.0	13.6	7.9	20.1	8.8	7.0	750
1980									
Jan.	4-8	13.0	11.9	6.2	20.0	10.7	11.4	19.8	720
	11-14	5.2	7.9	6.2	9.5	9.2	10.8	5.9	920
	14-17	14.0	31.0	17.6	15.8	12.6	14.8	15.0	680
	21-25	5.0	8.5	7.3	26.4	7.4	4.8	7.3	540
	25-29	25.2	11.4	19.3		67.8	10.1	9.2	680
	29-2	9.5	11.9	7.5		8.0	6.9	8.9	380
Feb.	2-6	18.6	7.6	11.5			9.9	10.0	840
mean		30.6	41.7	27.0	26.9	23.5	18.7	14.6	1147
(a + b + c) J			33.1			23.0		14.6	--
sediment, food			28.9			13.6		--	1000




Fig. 2. Relation between daily amount of food supplied and sediment. Upper and lower diagrams show the results obtained from sea bream and yellowtail farms, respectively.

Table 2. Seasonal variation of sediment flux ($dry-g/m^2 \cdot day$) in different farm.

Season	Sea bream St. 1 (8-12 m)	Yellowtail St. 2 (14-16 m)	mean
Summer (Jun-Jul)	15.7	12.1	13.9
Autumn (Sep-Nov)	44.6	28.8	36.7
Winter (Dec-Jan)	19.8	17.7	18.8

Table 3. The amount of sediment flux influenced by food quality.

Type of food	Date	SEDIMENT ($dry-g/m^2 \cdot day$)		FOOD (kg/day)	
		Sea bream (8-12 m)	Yellowtail (14-16 m)	Sea bream	Yellowtail
Frozen mackerel	(June 6- June 26)	21.6	17.0	1180	1314
Fresh anchovy	(June 27- July 13)	5.9	4.1	1353	1050

Bibliography

1. 森野静也 (1977): "浅海養殖と自家汚染", 31-41 (恒星社厚生閣, 東京).
 2. 花園 資 (1980): "赤潮に関する近年の知見と研究の問題点", 1~4 (日本水産資源保護協会, 東京).
 3. 半田暢彦 (1979): 対話「セジメント・トラップ」, 海洋科学, 11 (12), 959-964.
 4. 畑 幸彦 (1979): 海洋, 主として沿岸域における沈降粒子捕集器とその捕集物, 同上, 11 (12), 996-1002.
 5. 平田八郎・山口照男・植田隆史 (1979): 海面における溶存酸素量の水平的変化と海中懸濁物量との関連性, 琉球島周辺海域における陸棚漁場の開発利用に関する研究, 昭和53年度特定研究経過報告書, 44-53, 鹿大水.
 6. 井上裕雄・田中啓陽・福田 清 (1970): 浅海養魚場における海水の交流について II 田浦ハマチ養魚場の場合, 日本水産, 36, 776-782.
ISEKI, K., F. WHITNEY and C. S. WONG (1980): Biochemical Changes of Sedimented Matter in Sediment Trap in Shallow Coastal Waters. *Bull. Plankton Soc. Jap.*, 27 (1), 27-36.
 7. 門脇秀策・加世堂照男・中岡昌幸・平田八郎 (1978): 浅海養殖漁場における DO の航走連続記録 II 給餌に伴う浮遊懸濁物の拡散, 鹿大水紀要, 27, 281-288.
 8. 小山忠四郎 (1975): 湖沼堆植物の物質変化の機構に関する生物地球化学的考察, 水処理技術, 16, 19-39.
 9. 窪田敏文 (1977): "浅海養殖と自家汚染", 9-18 (恒星社厚生閣, 東京).
 10. 西沢 敏 (1966): 海中懸濁物: デトライタスから共棲小宇宙まで, 日本プランクトン学会報, 113, 1-33.
 11. 新田忠雄 (1980): "赤潮に関する近年の知見と問題点", 211-214 (日本水産資源保護協会, 東京).
 12. 田平紀男 (1978): "過密養殖防止のための漁場利用規制, ハマチ養殖に関する法的考察, 鹿児島県における赤潮の研究", 112-121 (鹿児島県漁業・他2団体, 鹿児島).
 13. 田中啓陽 (1977): "浅海養殖と自家汚染", 42-51 (恒星社厚生閣, 東京).
 14. 山内達也 (1979): シオミズツボウムシのフィードバックシステムにおける老廃物分解作用に関する研究, 鹿児島大学水産学部修士論文.
-
1. Shizuya HAGINO (1977): "Senkai Yoshoku to Jika Osen" (Shallow-waters Fish Farming and Auto-pollution) 31-41 (Kosei Co., Koseikaku, Tokyo).
 2. Tasuku* HANAOKA (1980): "Akashio ni kansuru Kinnen no Chiken to Kenkyu no Mondai" (Problems of views and studies about red tide), 1-4 (Nihon Suisan Shigen Hogo Kyokai/Japanese Association for the protection of marine resources. Tokyo).
 3. Nobuhiko HANDA (1979): "Taiwa "Sejimento - Torappu"" (Conversation "Sediment traps"). *Kaiyo Kagaku* (Marine Science), 11 (12), 959-964.
 4. Yukihiko* HATA (1979): "Kaiyo shu to shite Engan'iki ni okeru Chinko Ryushi Hoshuki to sono Hoshubutsu" (Devices to collect precipitated particles in the sea, particularly in coastal areas and the matters collected), ditto 11 (12), 996-1002.
 5. Hachiro HIRATA, Teruo YAMAGUCHI, Takashi MAKITA (1979): "Kaimen ni okeru Yozon Sansoryo no Suihei Henka to Kaichu Kendakubutsuryo to no Kanrensei / Ryukyu Toko Shuhen Kaiiki ni okeru Rikudana Gyojo no Kaihatsu Riyo ni kansuru Kenkyu" (Relationship between the horizontal variations of DO volumes at the surface of the sea and the volumes of suspended matters in the sea / Study about the

exploitation and use of the continental shelf in the area around the Ryukyu isles). Continuous report on specific research for the year 1978, 44-53, Kagoshima University, Fisheries Department.

6. Hiroo* INOUE, Keiyo TANAKA, Kiyoshi FUKUDA (1970): Senkai Yogyojo ni okeru Kaisui no Koryu ni tsuite - II Taura Hamachi Yogyojo no Baai (Water exchange in shallow-sea fish farms - II The yellowtail farms of Taura). Nihon Suisan Gakkaishi (Japanese Bulletin of the Scientific Fisheries), 36, 776-782.
7. Shusaku KADOWAKI, Teruo KASEDO, Tsurayuki NAKAZONO, Hachiro HIRATA (1978): Senkai Yoshoku Gyojo ni okeru DO no Koso Renzoku Kiroku - II. Kyuji ni tomonau Fuyu Kendakubutsu no Kakusan (Continuous recording of DO in coastal fish farms by cruising - II. The dispersion of sediments accompanying feeding). This journal, 27, 281-288.
8. Chushiro* KOYAMA (1975): Kosho Taiseikibutsu no Busshitsu Henka no Kiko ni kansuru Seibutsuchikyukagakuteki Kosatsu (Biogeochemical discussion about the changes in the substance of sediments in lakes and marshes). Mizushori Gijutsu (Water control technology), 16, 19-39.
9. Toshifumi KUBOTA (1977): Same references as in 1. above, 9-18.
10. Toshi NISHIZAWA (1966): Kaichu Kendakubutsu: Detoraitasu kara Kyosei Ko-Uchu made (Suspended matters in the sea: from detritus to the symbiotic microcosm). Nihon Purankuton Gakkai ho (Report of the Japanese Society for the Study of Plankton), 113, 1-33.
11. Nobuo NITTA (1980): Same reference as in 1. above, 211-214.
12. Norio TAHIRA (1978): Kamitsu Yoshoku Boshi no tame no Gyojo Riyo Kisei / Hamachi Yoshoku ni kansuru Hoteki Kosatsu. Kagoshima-wan ni okeru Akashio no Kenkyu (Regulations about the use of fishing grounds for the prevention of overdense fish farming / Legal discussion about yellowtail culture / Study on red tide in the bay of Kagoshima), 112-121 (Kagoshima Prefecture Fisheries Council and 2 other groups, Kagoshima).
13. Keiyo TANAKA (1977): Same reference as in 1. above, 42-51 (Kosei Co. Koseikaku, Tokyo).
14. Tatsya YAMAUCHI (1979): Shiomizu Tsubowamushi no Fidobakku Shisutemu ni okeru Rohaibutsu Bunkai Riyo ni kansuru Kenkyu (Study about the decomposition action of old waste in the feedback system of saltwater tsubo rotifers), Master's dissertation, Fisheries Department, Kagoshima University.

*Translator's note: Probable reading only; not quite certain.